

# Fire risk report for *Cenchrus clandestinus*

|                                                                                                                      |
|----------------------------------------------------------------------------------------------------------------------|
| <b>Full Species Name</b><br><i>Cenchrus clandestinus</i> (Hochst. ex Chiov.) Morrone                                 |
| <b>Family:</b> Poaceae                                                                                               |
| <b>Common names:</b><br>Kikuyu grass                                                                                 |
| <b>Synonyms:</b><br><i>Pennisetum clandestinum</i>                                                                   |
| Known occurrences (as of 2020)<br> |
| Year first documented as naturalized in Hawai'i: 1938                                                                |
| This species has been ranked by the Hawai'i Weed Risk Assessment program as High Risk with a score of 18.            |
| <a href="#">View photos on Starr Environmental</a>                                                                   |
| <a href="#">View on Wikipedia</a>                                                                                    |
| <a href="#">View occurrences on iNaturalist</a>                                                                      |
| <a href="#">View at Plants of Hawaii</a>                                                                             |
| <a href="#">View photos on Flickr</a>                                                                                |

0      **I**      .5      1  
Lowest risk      ⇔      Highest risk

This species is likely a **high** fire risk in Hawai'i with a fire risk score of **0.59**.

This species was ranked by 49 managers on a scale of 'no risk', 'low risk', 'medium risk', or 'high risk'. The numerical score ranges from 0 to 1 with higher scores indicating more managers considered it a higher risk. A score of > .31 indicates high risk.

| Summary of Fire ecology                       |         |
|-----------------------------------------------|---------|
| Native habitat fire proneness                 | No Data |
| Fire promoting plant in its native range      | No      |
| Fire promoting plant in its introduced range* | Yes     |
| Regenerates after fire                        | Yes     |
| Promoted by fire                              | No      |
| Reported flammable*                           | High    |
| Relative is flammable*                        | Yes     |

\*These values were used by the model to predict fire risk

## Detailed summary of Fire Ecology

|                                                                                                                                                                                                   |         |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Native habitat fire proneness (In any part of the plant's native range is its habitat described as fire prone due to natural or human caused fires?)                                              | No Data |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Fire promoting plant in its native range (Does the species act as a major fuel source, increase fire severity, frequency, or modify fuel bed characteristics within its native range?)            | No      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Fire promoting plant in its introduced range (Same as Fire Promoting Native but within the species introduced range)                                                                              | Yes     | <p>"ungrazed kikuyu could carry catastrophic fires (flame lengths 3.0 m, covering &gt;75 ha after one hour)."</p> <p>Blackmore, M., &amp; Vitousek, P. M. (2000). Cattle Grazing, Forest Loss, and Fuel Loading in a Dry Forest Ecosystem at Pu'u Wa'aWa'a Ranch, Hawai'i 1. <i>Biotropica</i>, 32(4a), 625-632.</p> <p>-----</p> <p>"Kikuyu grass creates a considerable fire hazard as the old stolons die and accumulate because of the new growth scrambling over the top" (Edgar &amp; Connor, 2000; p. 574)"</p> <p><a href="http://www.hear.org/pier/species/pennisetum_clandestinum.htm">http://www.hear.org/pier/species/pennisetum_clandestinum.htm</a></p> |
| Regenerates after fire (Does the plant regrow after fire by any means? This includes resprouters, reseeder, and recruiters which dispersed into the area within approximately one year post fire) | Yes     | <p>"Burning significantly reduced the aboveground biomass of residual material from 3.54 to 1.40 t/ha [implies regeneration]"</p> <p>Holliday, J., du Toit, J. C. O., &amp; Morris, C. D. THE EFFECT OF BURNING ON ABOVEGROUND BIOMASS AND CHEMICAL COMPOSITION ON PENNISETUM CLANDESTINUM-DOMINATED GRASSLAND. <i>Weed research</i> 36, 213-225</p> <p>-----</p> <p>"According to older fire users, this grass has existed in the region for ,20–40 years but became an agricultural pest ,10–15 years ago and that despite burning (or repeated</p>                                                                                                                 |

|                                                                                                                   |      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|-------------------------------------------------------------------------------------------------------------------|------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                                                                   |      | <p>burning) and the use of herbicides it always reinvades agricultural plots (Fig. 2"</p> <p>Martínez-Torres, H. L., Castillo, A., Ramírez, M. I., &amp; Pérez-Salicrup, D. R. (2016). The importance of the traditional fire knowledge system in a subtropical montane socio-ecosystem in a protected natural area. <i>International Journal of Wildland Fire</i>, 25(9), 911-921.</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Promoted by fire (Does the plant increase in abundance after a fire?)                                             | No   |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| Reported flammable (Is the species described as being flammable, being a major wildfire fuel, or high fire risk?) | High | <p>"ungrazed kikuyu could carry catastrophic fires (flame lengths 3.0 m, covering &gt;75 ha after one hour). "</p> <p>Blackmore, M., &amp; Vitousek, P. M. (2000). Cattle Grazing, Forest Loss, and Fuel Loading in a Dry Forest Ecosystem at Pu'u Wa'aWa'a Ranch, Hawai'i 1. <i>Biotropica</i>, 32(4a), 625-632.</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Relative is flammable (Does a plant in the same genus meet the Reported Flammable criteria?)                      | Yes  | <p>"Fuels: Buffelgrass [<i>Cenchrus ciliaris</i>] fine fuel loads are generally much higher than fine fuel loads from native plants in desert environments. Thus, fires in buffelgrass stands may have longer flame lengths, greater rates of spread, and higher temperatures than fires in native desert vegetation, and cause high mortality in native flora and fauna [43]. Buffelgrass stands burn "very hot" [24] and can burn when green [42,129]. In the Sonoran Desert, buffelgrass-fueled fires can reach temperatures so hot that the soil is scorched and the bedrock cracked [42]. Headfires in buffelgrass stands can reach temperatures of 1,090 to 1,300 °F (585-700 °C) [27,103]. Esque and others [42] state that buffelgrass grows into an "almost-woody subshrub", accumulating flammable material over several years, "in effect unlinking fire frequency from annual climatic variability and increasing the fire intensity". "</p> <p><a href="https://www.fs.fed.us/database/feis/plants/graminoid/pencil/all.html#FIRE%20ECOLOGY">https://www.fs.fed.us/database/feis/plants/graminoid/pencil/all.html#FIRE%20ECOLOGY</a></p> <p>-----</p> <p>"Buffel grass invasion was significantly correlated with increased fuel loads. Increased fuel loads were significantly correlated with increased burn severity, although the direct relationship between the proportion of buffel grass and increased burn severity was marginally non-significant."</p> |

|  |  |                                                                                                                                                                                                                                                                 |
|--|--|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|  |  | Miller, G., Friedel, M., Adam, P., & Chewings, V. (2010). Ecological impacts of buffel grass ( <i>Cenchrus ciliaris</i> L.) invasion in central Australia—does field evidence support a fire-invasion feedback?. <i>The Rangeland Journal</i> , 32(4), 353-365. |
|--|--|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

Text in quotes are direct quotes from the source

Text in square brackets was added by the assessor to clarify something or to summarize from a figure.

Text preceded by a “#” is comment from the assessor

---

The data presented were assembled from literature and database searches for each species using as much data as could be collected regarding the plant’s fire ecology under natural conditions. Searches aimed to be exhaustive and consist of as much data as could be located in 2020. Our machine learning algorithm was trained on 49 species of plants which had their fire risk ranked by 49 managers in Hawai’i in November 2020. The model used a conditional random forest regression algorithm to predict scores for each species using the manager score as the response variable and the fire ecology traits of each plant as the predictor variables to generate a fire risk score. This trained model was then used to predict the fire risk for all species which were not ranked by managers. The model was calibrated such that it is 90% accurate at predicting high fire risk plants and 79% accurate at predicting low fire risk plants. This research and the resulting fire risk model has been published in the journal [Biological Invasions](#) by [Kevin Faccenda](#) and [Curt Daehler](#) (both at the University of Hawai’i at Mānoa).

Note that the analysis doesn’t account for a plant species’ spatial distribution, population density, or distinct climate and ecosystem conditions (which can also influence fire risk). The fire risk of these species are mostly under “worst case” environmental conditions where the climate is dry enough to maintain fire, but wet enough to allow for plant growth and fuel accumulation. The fire risk ranking should not be taken as a stand-alone risk metric in prioritizing weed control efforts. Rather, this information may also be useful for determining if a newly discovered species poses a potential fire threat in wildland areas.

More general information on the weed risks and ecology of non-native plants in Hawai’i is available from the Hawai’i Invasive Species Committee’s [Weed Risk Assessment database](#).

View more fact sheets at <https://www.pacificfireexchange.org/weed-fire-risk-assessments>

---

Fact sheet prepared by Kevin Faccenda ([faccenda@hawaii.edu](mailto:faccenda@hawaii.edu)) in November 2021. Data were prepared by Kevin Faccenda in 2020.

This research was funded by the Department of the Interior Pacific Islands Climate Adaptation Science Center. The project described in this publication was supported by Grant or Cooperative Agreement No.G20AC00073 to Curt Daehler from the United States Geological Survey. The views and conclusions contained in this document are those of the authors and should not be interpreted as representing the opinions or policies of the U.S. Geological Survey. Mention of trade names or commercial products does not constitute their endorsement by the Pacific Islands Climate Adaptation Science Center or the National Climate Adaptation Science Center or the US Geological Survey.

