


## Fire risk report for *Parthenium hysterophorus*

<b>Full Species Name</b> <i>Parthenium hysterophorus</i> L.
<b>Family:</b> Asteraceae
<b>Common names:</b> false ragweed Santa Maria
<b>Synonyms:</b>
Known occurrences (as of 2020) 
Year first documented as naturalized in Hawai'i: 1960
This species has not yet been ranked by the Hawai'i Weed Risk Assessment program as of 2020.
<a href="#">View photos on Starr Environmental</a>
<a href="#">View on Wikipedia</a>
<a href="#">View occurrences on iNaturalist</a>
<a href="#">View at Plants of Hawaii</a>
<a href="#">View photos on Flickr</a>

0                      .5                      1  
Lowest risk                      Highest risk

This species is likely a **low** fire risk in Hawai'i with a fire risk score of **0.25**.

This species was ranked by our machine learning algorithm using the data presented on the next page. A predicted score of > .34 suggests the plant is a high fire risk.

Summary of Fire ecology	
Native habitat fire proneness	Uncertain
Fire promoting plant in its native range	No
Fire promoting plant in its introduced range*	No
Regenerates after fire	Yes
Promoted by fire	Yes
Reported flammable*	Low
Relative is flammable*	No

\*These values were used by the model to predict fire risk

## Detailed summary of Fire Ecology

Native habitat fire proneness (In any part of the plant's native range is its habitat described as fire prone due to natural or human caused fires?)	Uncertain	
Fire promoting plant in its native range (Does the species act as a major fuel source, increase fire severity, frequency, or modify fuel bed characteristics within its native range?)	No	
Fire promoting plant in its introduced range (Same as Fire Promoting Native but within the species introduced range)	No	<p>"Parthenium dominant pastures will not carry a fire. Therefore a reasonable amount of grass is necessary to provide fuel for an effective burn to occur."</p> <p><a href="https://www.agrifutures.com.au/wp-content/uploads/publications/06-130.pdf">https://www.agrifutures.com.au/wp-content/uploads/publications/06-130.pdf</a></p> <p>W Vogler, S Navie, S Adkins and C Setter. 2006. Use of Fire to Control Parthenium Weed. A report for the Rural Industries Research and Development Corporation</p>
Regenerates after fire (Does the plant regrow after fire by any means? This includes resprouters, reseeder, and recruiters which dispersed into the area within approximately one year post fire)	Yes	<p>"Fire has been studied in Australia as a measure to control parthenium weed infestations. The results have shown that the overall effect is to create open niches in the landscape, in which a larger number of parthenium weed seed are able to germinate in the absence of vegetation.<sup>56</sup> Therefore, management of parthenium weed in pastures through burning is not considered to be an option."</p> <p><a href="https://doi.org/10.1002/ps.3708">https://doi.org/10.1002/ps.3708</a></p> <p>Adkins, S., &amp; Shabbir, A. (2014). Biology, ecology and management of the invasive parthenium weed (<i>Parthenium hysterophorus</i> L.). <i>Pest management science</i>, 70(7), 1023-1029.</p>
Promoted by fire (Does the plant increase in abundance after a fire?)	Yes	<p>"Fire has been studied in Australia as a measure to control parthenium weed infestations. The results have shown that the overall effect is to create open niches in the landscape, in which a larger number of parthenium weed seed are able</p>

		<p>to germinate in the absence of vegetation.<sup>56</sup> Therefore, management of parthenium weed in pastures through burning is not considered to be an option."</p> <p><a href="https://doi.org/10.1002/ps.3708">https://doi.org/10.1002/ps.3708</a></p> <p>Adkins, S., &amp; Shabbir, A. (2014). Biology, ecology and management of the invasive parthenium weed (<i>Parthenium hysterophorus</i> L.). <i>Pest management science</i>, 70(7), 1023-1029.</p> <p>-----</p> <p>"There was an increase in parthenium frequency after the initial spring and autumn burns, with a subsequent large decline in future years even though follow-up burns were conducted. In the ungrazed control treatments parthenium frequency declined in the first year and remained at a low"</p> <p><a href="https://www.agrifutures.com.au/wp-content/uploads/publications/06-130.pdf">https://www.agrifutures.com.au/wp-content/uploads/publications/06-130.pdf</a></p> <p>W Vogler, S Navie, S Adkins and C Setter. 2006. Use of Fire to Control Parthenium Weed. A report for the Rural Industries Research and Development Corporation</p>
Reported flammable (Is the species described as being flammable, being a major wildfire fuel, or high fire risk?)	Low	<p>"On rangeland the burning of parthenium weed by wild-fire followed by the planting of a vigorous pasture has been successful in trials and is proposed as the most practical method of control of the weed under these conditions. Unfortunately, the burning of dense stands of this weed is not possible where other vegetation is not present or until the remains of two or more season's growth are present. The aerial application of low rates of a selective herbicide such as 2.4-D is also proposed to allow the grass to compete more successfully."</p> <p>Haseler, W. H. (1976). <i>Parthenium hysterophorus</i> L. in Australia. <i>PANS</i>, 22(4), 515–517.</p> <p><a href="https://doi.org/10.1080/09670877609414342">https://doi.org/10.1080/09670877609414342</a></p> <p>-----</p> <p>"Parthenium dominant pastures will not carry a fire. Therefore a reasonable amount of grass is necessary to provide fuel for an effective burn to occur."</p> <p><a href="https://www.agrifutures.com.au/wp-content/uploads/publications/06-130.pdf">https://www.agrifutures.com.au/wp-content/uploads/publications/06-130.pdf</a></p> <p>W Vogler, S Navie, S Adkins and C Setter. 2006. Use of Fire to Control Parthenium Weed. A report for the Rural Industries Research and Development Corporation</p>
Relative is flammable (Does a plant in the same genus meet the	No	

Reported Flammable criteria?)		
-------------------------------	--	--

Text in quotes are direct quotes from the source

Text in square brackets was added by the assessor to clarify something or to summarize from a figure.

Text preceded by a “#” is comment from the assessor

---

The data presented were assembled from literature and database searches for each species using as much data as could be collected regarding the plant’s fire ecology under natural conditions. Searches aimed to be exhaustive and consist of as much data as could be located in 2020. Our machine learning algorithm was trained on 49 species of plants which had their fire risk ranked by 49 managers in Hawai’i in November 2020. The model used a conditional random forest regression algorithm to predict scores for each species using the manager score as the response variable and the fire ecology traits of each plant as the predictor variables to generate a fire risk score. This trained model was then used to predict the fire risk for all species which were not ranked by managers. The model was calibrated such that it is 90% accurate at predicting high fire risk plants and 79% accurate at predicting low fire risk plants. This research and the resulting fire risk model has been published in the journal [Biological Invasions](#) by [Kevin Faccenda](#) and [Curt Daehler](#) (both at the University of Hawai’i at Mānoa).

Note that the analysis doesn’t account for a plant species’ spatial distribution, population density, or distinct climate and ecosystem conditions (which can also influence fire risk). The fire risk of these species are mostly under “worst case” environmental conditions where the climate is dry enough to maintain fire, but wet enough to allow for plant growth and fuel accumulation. The fire risk ranking should not be taken as a stand-alone risk metric in prioritizing weed control efforts. Rather, this information may also be useful for determining if a newly discovered species poses a potential fire threat in wildland areas.

More general information on the weed risks and ecology of non-native plants in Hawai’i is available from the Hawai’i Invasive Species Committee’s [Weed Risk Assessment database](#).

View more fact sheets at <https://www.pacificfireexchange.org/weed-fire-risk-assessments>

---

Fact sheet prepared by Kevin Faccenda ([faccenda@hawaii.edu](mailto:faccenda@hawaii.edu)) in November 2021. Data were prepared by Kevin Faccenda in 2020.

This research was funded by the Department of the Interior Pacific Islands Climate Adaptation Science Center. The project described in this publication was supported by Grant or Cooperative Agreement No.G20AC00073 to Curt Daehler from the United States Geological Survey. The views and conclusions contained in this document are those of the authors and should not be interpreted as representing the opinions or policies of the U.S. Geological Survey. Mention of trade names or commercial products does not constitute their endorsement by the Pacific Islands Climate Adaptation Science Center or the National Climate Adaptation Science Center or the US Geological Survey.

